

Original Article

Regulation of autonomic functions following two high frequency yogic breathing techniques

Joydeb Mondal¹, Ragavendrasamy Balakrishnan², Manjunath Nandi Krishnamurthy^{2,*}

¹S-VYASA Yoga University, Bangalore, India; ²Department of Life Sciences, S-VYASA University, Bangalore, India

ABSTRACT

Yoga is an ancient Indian system of life, encompassing various practices including practices for self-discipline and also for regulating the health states of the individual, being practiced for thousands of years. The present study aims at understanding the effect of two high frequency breathing practices over autonomic nervous system. Forty healthy male volunteers of age 21 ± 2 years with 9 ± 3 months of *Yoga* practice experience were recruited. The two high frequency *Yoga* breathing practices, *kapalabhati* (KB) and *bhastrika* (BH) were given as interventions randomly on either of the two days to minimise laboratory bias. They were assessed before and immediately after the interventions for heart rate, respiratory rate, heart rate variability (HRV), blood pressure and peripheral oxygen saturation. There was a significant increase in heart rate ($p < 0.01$; $p < 0.001$), systolic blood pressure ($p < 0.01$; $p < 0.001$), NN50 ($p < 0.01$; $p < 0.001$) component of HRV for both KB and BH groups respectively. There was a significant reduction in respiratory rate in both the groups ($p < 0.001$, and $p < 0.05$, BH and KB respectively) immediately following intervention. A significant increase in LF component of HRV and reduction in Diastolic blood pressure and high frequency (HF) component following KB was also observed ($p < 0.05$, for all comparisons). The Mean peripheral oxygen saturation remained unaltered in both the groups ($p > 0.05$). The results suggest that high frequency *yoga* breathing practices induce physiological arousal immediately as evidenced by increased blood pressure and heart rate. The sympathetic arousal was more following KB session as evidenced by an increased diastolic blood pressure, LF power and a decrease in HF power of HRV as compared to the BH session.

Keywords pranayama, kapalabhati, bhastrika, autonomic functions, heart rate variability

INTRODUCTION

Yoga is a way of life being practiced in India since thousands of years in the form of physical postures (*asana*), breathing practices (*pranayama*), and meditation (*dhyana*). Past decades have encouraged rigorous scientific studies to understand the science behind these traditional practices towards understanding their possible role in promoting health and alleviate disease conditions. Recent studies in *Yoga* have demonstrated beneficial effects in many cardio vascular diseases and metabolic disorders like diabetes mellitus (Ren et al., 2012).

Present day science has started recognizing the relationship between the emotional regulation, breathing and metabolism (Porges, 2001), which has been described in the *hatha yoga pradipika* (Muktibodhananda, 2000). Also, respiratory patterns and maneuvers have shown to provide striking influences on the autonomic nervous system and may exacerbate or reduce adverse responses to stressors. For example, increased breathing rate is a typical response to stressful situations (Grossman, 1983). This tendency can lead to breathing in

excess of metabolic needs (hyperventilation), which causes reduction in blood carbon dioxide concentrations. The reduced carbon dioxide causes psycho-physiological and psychological effects that include (a) enhanced arousal and anxiety and (b) decreased cerebral and coronary blood flow, which can lead to a variety of clinical symptoms including dizziness, poor performance, headache, chest pain, cardiac abnormalities, and sleep disturbances (Brown, 1953; Fried, 1988; Grossman, 1983; Lamb et al., 1958; Magarian, 1982). Certain other respiratory patterns that modestly elevate blood carbon dioxide concentration appear to promote the opposite effects, including reduced anxiety and increased or well-maintained cerebral and coronary blood flow (Grossman, 1983). Reduction of hypertension (Irvine et al., 1986; Patel et al., 1981; Patel and North, 1975) and dramatic improvement of heart disease have resulted from integrated treatment programs that included *yogic* breathing practices (Berman and Larson, 1994).

Pranayama, the science and art of breath regulation is more than a simple breathing exercise. It is not merely breathe control but one of the powerful yogic techniques used to regulate the *prana* (defined as the force behind every action) in the body to a higher frequency (Singh et al., 1990). *Pranayama* techniques involve breathing through either of the nostrils or through both the nostrils, and conscious regulation of rate and depth of breathing (Niranjanananda, 2000).

Kapalabhati (KB) is mentioned as one of the six cleansing procedures (Muktibodhananda, 2000) that involves rapid breathing consisting of active expiration with the help of abdominal muscles and passive spontaneous inspiration

*Correspondence: Manjunath Nandi Krishnamurthy

E-mail: nkmsarma@svyasa.org

Received April 7, 2014; Accepted February 12, 2015; Published February 28, 2015

doi: <http://dx.doi.org/10.5667/tang.2014.0015>

© 2015 by Association of Humanitas Medicine

This is an open access article under the CC BY-NC license.

(<http://creativecommons.org/licenses/by-nc/3.0/>)

happening during relaxation (Raghuraj et al., 1998). *Bhastrika* (BH) is another high frequency breathing practice similar to that of KB except for the forceful inhalation unlike KB (Muktibodhananda, 2000).

However, there being very subtle difference in the actual procedure of the practice, these practices have been described as voluntary breath regulation practices and cleansing procedures thereby demanding closer observation into the traditional literatures to unearth their significance and understand their possible applications.

Experimental data shows that KB and BH practices have a direct influence on physiological processes such as respiratory and cardiovascular system, biochemical parameters (Desai and Gharote, 1990) and central nervous activity (Stancák et al., 1991). An increase in the performance in a letter cancellation task was observed immediately after practice of KB at a rate of 60 strokes (breaths) per minute (Telles et al., 2008). Whereas the cognitive evoked potentials showed a reduction in latency and increase in amplitude following practice of KB when administered at the rate of 120 strokes (breaths) per minute (Joshi and Telles, 2009). Earlier studies have reported that slower breathing practices, alternate nostril breathing (*nadishuddhi*) increase parasympathetic activity whereas faster breathing practices (KB) increase sympathetic activity (Pal et al., 2004).

Heart Rate Variability (HRV) is an indicator of the autonomic nervous system activity of an individual. Previous studies conducted on the influence of *kapalabhati* on HRV have shown increased sympathetic arousal (Raghuraj et al., 1998). However studies are lacking in understanding the difference in autonomic changes following BH and KB. BH and KB both being high frequency breathing practices it is a mandate to understand the underlying difference in their mechanism of action. This preliminary study is expected to aid in understanding the commonalities and differences between these two high frequency *yoga* breathing techniques on components of autonomic functions.

Methodology

Subjects

An advertisement mentioning 'volunteers shall register for a study to understand the autonomic regulations of two breathing practices' was made in a residential *yoga* institute campus. Of the sixty registered volunteers, 40 volunteers of the age group (21 ± 2 years), having an experience of 9 ± 3 months, who were clinically healthy, not under medication from the date of start of study, were recruited for the study following approval from the Institutional Ethics Committee. A written informed consent was obtained from all the subjects before recruitment.

Subjects were assessed five minutes before and immediately after KB on one day and BH on another day, but at the same time of the day with similar physical mental load before coming to the laboratory. Alternate allocation of KB and BH on day one of their visit to the lab was done to ensure 50% of the subjects performing KB on Day one while the remaining 50% performed BH.

Immediately after reaching the laboratory, subjects were made to sit in a sound attenuated, dim light room, allowing them to relax for five minutes, following which baseline recordings of blood pressure, heart rate, ECG, oxygen saturation, and respiration were recorded. The intervention KB and BH were randomly administered for duration of 1 min.

Assessments

Heart Rate Variability

The ECG was recorded using a polygraph (BIOPAC MP100, USA) at a sampling rate 1024 Hz and was analyzed offline. The data were acquired using the standard limb lead II configuration in five minutes epochs before and immediately after the intervention. The data was visually inspected offline and noise free data were included for analysis. The R data waves were detected to obtain a point event series of successive RR intervals, from which the beat to beat heart series were computed. The data were analysed with an HRV analysis program, 'Kubios', developed by the biomedical signal analysis group (Finland).

Respiration (R)

Respiration was as recorded using a stetho-graph fastened approximately 1 cm below the margin of the rib cage while subject was made to sit erect. Respiration was constantly monitored before and immediately following the intervention for 5 mins.

Blood pressure (BP)

A digital sphygmomanometer device (Mindray, Hong Kong) was used to monitor the blood pressure. The digital manometer was tied in the left arm over the brachial artery and was inflated to record the blood pressure immediately before the start and after the completion of the intervention.

Oxygen Saturation

Oxygen saturation was measured using Pulse oxymeter (Mindray, Hong Kong). A photo plethysmograph comprising of an infra-red diode was connected to the middle finger of the right hand and continuous blood oxygen saturation was recorded before and after the intervention for five minutes.

Intervention

The participants were made to sit in a chair having their spine straight and the hands resting on the knees. With their eyes closed, they were asked to voluntarily relax the entire body. Ag-AgCl electrodes were placed as per limb lead II and respiration belt was applied on the chest 1 cm below the lower margin of the rib cage. Sphygmomanometer cuff was tied to the left arm and the Pulse Oximeter was fixed to the thumb of the right hand to avoid any interference from the sphygmomanometer cuff.

KB and BH was administered as mentioned in *Hatha Yoga Pradipika* (Mukthibodhananda, 2000).

KB: The participants were asked to sit in a chair having the spine straight and the hands resting on the knees. The subjects were instructed to relax the whole body. Inhaling deeply through both nostrils, expanding the abdomen, the subject has to exhale with a forceful contraction of the abdominal muscles. Following forceful exhalation, voluntary relaxation of the abdominal muscles promotes downward movement of the diaphragm causing a negative pressure in the thoracic cavity, facilitating rapid influx of air. The speed of the practice was guided with the help of an electronic timer device. After completion of 60 rapid breaths in succession, the participants were asked to relax in the same position without voluntarily manipulating their breath. (Muktibodhananda, 2000).

BH: The subjects were instructed to breathe in and out forcefully through the nose without straining any part of the body. Inhaling deeply through both the nostrils forcefully, the subjects were instructed to exhale forcefully for 40 rounds in a rhythmic fashion. The speed of the practice was guided with

Table 1. The group mean \pm SD of HRV, SpO₂, RR and BP recorded before and immediately after the practice of *Kapalabhati* and *Bhastrika Pranayama*

VARIABLES	BHASTRIKA		KAPALABHATI	
	Pre	Post	Pre	Post
Mean HR (bpm)	80.05 \pm 10.53	83.28 \pm 9.17***	80.59 \pm 14.78	82.46 \pm 14.52***
NN50 (nu)	87.87 \pm 42.73	76.28 \pm 66.60***	88.49 \pm 13.70	80.05 \pm 15.95***
LF (nu)	54.20 \pm 16.88	57.98 \pm 19.23	50.27 \pm 17.87	57.75 \pm 18.87*
HF (nu)	45.78 \pm 16.89	42.00 \pm 19.23	49.72 \pm 17.87	42.22 \pm 18.87*
LF/HF RATIO (nu)	1.65 \pm 1.43	1.99 \pm 1.55	1.41 \pm 1.35	1.90 \pm 1.78
RR (bpm)	15.29 \pm 2.40	13.82 \pm 2.20***	15.56 \pm 2.31	14.85 \pm 2.49*
SpO ₂ (%)	96.97 \pm 0.75	96.82 \pm 4.75	97.10 \pm 0.97	96.97 \pm 0.95
BP-SYS (mmHg)	107.66 \pm 8.4	110.35 \pm 11.48***	107.40 \pm 8.70	111.41 \pm 10.88**
BP-DIA (mmHg)	72.56 \pm 5.70	72.42 \pm 13.42	73.23 \pm 7.60	76.09 \pm 8.46*

Paired 't' test indicating within group differences * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$. Mean HR (bpm): mean heart rate in beats per minute, NN50 (nu): number of successive RR intervals differing more than 50 ms represented in normalised units, LF (nu): low frequency in normalised units, HF (nu): high frequency in normalised units, LF/HF Ratio (nu): ratio of low frequency and high frequency components in normalised units, RR (bpm): respiratory rate in normalised units, SpO₂ (%): percentage of oxygen in peripheral capillaries, BP-SYS: systolic blood pressure, BP-DIA: diastolic blood pressure.

the help of an electronic timer device. (Muktibodhananda, 2000).

Statistical analysis

Statistical analysis was performed using the SPSS version 19.0. Following the normal distribution of the data as indicated through Shapiro-wilk's test, within group differences were analyzed using paired 't' test and between group effects were analyzed using independent 't' test.

RESULTS

Paired 't' test performed (Table 1) to understand the within group differences showed a significant increase in heart rate ($p < 0.001$), and a significant decrease in NN50 ($p < 0.001$) immediately after the practice of BH and KB practices. A significant increase in LF ($p < 0.05$) and significant decrease in HF ($p < 0.05$) components of HRV was observed immediately following the practice of KB. There were no significant differences following intervention on both the days.

Respiratory rate was significantly reduced immediately following both the practices KB ($p < 0.05$) and BH ($p < 0.001$) (Table 1). There was no significant change observed in the peripheral oxygen saturation levels.

DISCUSSION

The purpose of this study was to understand the influence of two High Frequency *Yogic* Breathing practices on the Heart Rate Variability and Blood Pressure in healthy experienced *Yoga* practitioners. Both KB and BH practices involve constant forceful rapid exhalations; the difference in the two practices being additional active forceful inhalation in BH practice, which is thought to contribute to additional load over the physiological reserve. However, there were no studies comparing the role of KB and BH practices regulating the Heart Rate Variability.

The findings of the present study are in accordance with the

earlier findings indicating an increase in Sympathetic Nervous System activity immediately after both the practices. However, the sympathetic arousal was non-significantly more following the practice of BH as compared to KB, which can be attributed to the increased effort required to perform the BH, requiring more Cardiac output to suffice the needs of the exercising respiratory musculature and the diaphragm.

High frequency breathing is known to cause a direct influence on physiological processes through the cardio respiratory system (Stancák et al., 1991).

Though KB was shown to be a stimulating practice, reported to improve attention and concentration (Tells et al., 2008), there was a need to understand simpler breathing based methods to increase sympathetic activity. It is clear from the present study that high frequency breathing has a direct impact on the autonomic nervous system. While BH showed more influence on blood pressure and heart rate, KB had better influence on heart rate variability as expected. BH practice appears to influence the cardio Respiratory system more mechanically than KB.

The thoraco-abdominal manoeuvring involved in BH results in increased breath volume and greater contraction and expansion of the thorax and abdominal wall. In comparison, KB with more of diaphragmatic movement appears to exert less mechanical pressure on the thorax and hence has less influence on the blood pressure. No changes observed in oxygen saturation levels can probably be attributed to the duration of the practice.

The findings of the present study are in accordance with the earlier findings of high frequency breathing practices directly regulating the autonomic nervous system. The results suggest that practice of *Kapalabhati* and *Bhastrika* for 60 and 40 strokes per minute respectively in experienced practitioners, cause an increase in heart rate, blood pressure, LF component of HRV and a significant decrease in NN50 component and respiratory rate. Further follow up studies are required to understand the long term autonomic regulation following breathing practices. And evaluation of safety in hypertensive patients is warranted.

ACKNOWLEDGEMENTS

None.

CONFLICT OF INTEREST

The authors have no conflicting financial interests.

REFERENCES

- Berman BM, Larson DB. *Alternative medicine: Expanding medical horizons*. 1st ed. (Washington DC, North America: Diane Publishing Company), 1994.
- Brown EB. Physiological effects of hyperventilation. *Physiol Rev*. 1953;33: 445-471.
- Desai BP, Gharote ML. Effect of Kapalabhati on blood urea, creatinine and tyrosine. *Act Nerv Super (Praha)*. 1990;32:95-98.
- Fried R. The hyperventilation syndrome--research and clinical treatment. *J Neurol Neurosurg Psychiatry*. 1988;51:1600-1601.
- Grossman P. Respiration, stress, and cardiovascular function. *Psychophysiology*. 1983;20:284-300.
- Irvine MJ, Johnston DW, Jenner DA, Marie GV. Relaxation and stress management in the treatment of essential hypertension. *J Psychosom Res*. 1986;30:437-450.
- Joshi M, Telles S. A nonrandomized non-naive comparative study of the effects of kapalabhati and breath awareness on event-related potentials in trained yoga practitioners. *J Altern Complement Med*. 2009;15:281-285.
- Lamb LE, Dermksian G, Sarnoff C. Significant cardiac arrhythmias induced by common respiratory maneuvers. *Am J Cardiol*. 1958;2:563-571.
- Magarian GJ. Hyperventilation Syndromes: infrequently recognized common expressions of anxiety and stress. *Medicine (Baltimore)*. 1982;61:219-236.
- Muktibodhananda S. *Hatha Yoga Pradipika*. 2nd ed. (Bihar, India: Nesma Books India), 2000.
- Niranjananda S. *Prana, Pranayama Pranavidya*. (Jharkhand, India: Satyanandashram), 2000.
- Pal GK, Velkumary S, Madanmohan. Effect of short-term practice of breathing exercises on autonomic functions in normal human volunteers. *Indian J Med Res*. 2004;120:115-121.
- Patel C, North WR. Randomised controlled trial of yoga and bio-feedback in management of hypertension. *Lancet*, 1975;2:93-95.
- Patel C, Marmot MG, Terry DJ. Controlled trial of biofeedback-aided behavioural methods in reducing mild hypertension. *Br Med J (Clin Res Ed)*. 1981;282:2005-2008.
- Porges SW. The polyvagal theory: phylogenetic substrates of a social nervous system. *Int J Psychophysiol*. 2001;42:123-146.
- Raghuraj P, Ramakrishnan AG, Nagendra HR, Telles S. Effect of two selected yogic breathing techniques on heart rate variability. *Indian J Physiol Pharmacol*. 1998;42:467-472.
- Ren P, Barreto A, Gao Y, Adjouadi M. Comparison of the use of pupil diameter and galvanic skin response signals for affective assessment of computer users. *Biomed Sci Instrum*. 2012;48:345-350.
- Singh V, Wisniewski A, Britton J, Tattersfield A. Effect of yoga breathing exercises (pranayama) on airway reactivity in subjects with asthma. *Lancet*, 1990;335:1381-1383.
- Stancák A, Kuna M, Srinivasan, Dostálek C, Vishnudevananda S. Kapalabhati--yogic cleansing exercise. II. EEG topography analysis. *Homeost Health Dis*. 1991;33:182-189.
- Telles S, Raghuraj P, Arankalle D, Naveen KV. Immediate effect of high-frequency yoga breathing on attention. *Indian J Med Sci*. 2008;62:20-22.