

Contemporary Approach to Autotransplantation of Teeth with Complete Roots Using 3D-printing Technology

Jungha Park¹, Sangho Lee¹, Nanyoung Lee¹, Myoungkwan Jih¹, Hyeran Cheong²

¹*Department of Pediatric Dentistry, School of Dentistry, Chosun University*

²*Department of Dental Hygiene, Dong-A College of Health*

Abstract

Autotransplantation of teeth with complete roots is being increasingly performed for orthodontic treatment or for replacement of unsalvageable teeth, but this procedure has lower survival and success rates than those obtained following transplantation of teeth with incomplete root formation.

While previous autotransplantation procedures used only models of the donor tooth and recipient site, surgical guide templates created using 3D-printing technology are now available. They allow shaping of the recipient site in proper direction and to the correct depth, thereby reducing the treatment time and improving the success and survival rates.

Herein, we report a case involving autotransplantation of a tooth with complete roots at the site of a congenitally missing permanent tooth using surgical guide templates generated with 3D-printing technology. The procedure resulted in favorable healing of the transplanted tooth.

Key words : Autotransplantation, Teeth with complete roots, 3D-printing technology, Surgical guide templates

I. Introduction

Autotransplantation of teeth with complete roots has lower survival and success rates compared with those for teeth with incomplete roots[1,2]. Nevertheless, autotransplantation of teeth with complete roots is performed as part of orthodontic treatments and to replace unsalvageable teeth[3].

Ideally, root development in a tooth intended for successful autotransplantation should be half to two-thirds complete[4]. In addition, Hertwig's epithelial root sheath should be conserved at the 4th or 5th phase of root development as determined by the Moorrees' staging method[5]. Since the periapical regions of teeth with incomplete roots are covered by a

thin follicle or periodontal ligaments, only minimal force is required to remove such teeth from the alveolar socket. Therefore, minimal trauma is exerted on the periodontal ligaments during extraction. Furthermore, as blood supply is possible through the large apical foramen, there should be relatively good pulp healing[6]. In contrast, teeth with complete roots are at high risk of periodontal ligament damage during extraction, and exhibit relatively lower success and survival rates[6].

The reported success and survival rates of autotransplanted teeth with complete root formation vary substantially. In a study with a mean follow-up period of 63.8 months, Yoshino *et al.*[7,8] reported survival rates of 90.1% at 5 years and 70.5% at 10 years for 614 transplanted teeth. In a study of 38

Corresponding author : Sangho Lee

Department of Pediatric Dentistry, College of Dentistry, Chosun University, 303 Pilmun-daero, Dong-gu, Gwangju, 61452, Republic of Korea

Tel: +82-62-220-3860 / Fax: +82-62-225-8240 / E-mail: shclee@chosun.ac.kr

Received October 24, 2016 / Revised December 27, 2016 / Accepted December 12, 2017

www.kci.go.kr

transplanted teeth with a mean follow-up period of 9.2 years, Watanabe *et al.*[2,8] reported a survival rate of 86.8% and a success rate of 63.1%. In a study of 33 transplanted teeth with a mean follow-up period of 26.4 years, Czochrowska *et al.*[7,9] reported survival and success rates of 90% and 79%, respectively. Although the reported survival and success rates of autotransplantation of teeth with complete roots are diverse, they are lower than those of immature teeth. Therefore, studies that focus on the success rate of autotransplantation of teeth with complete roots are significant.

Factors influencing the prognosis of autotransplantation include anatomical and morphological characteristics of the donor tooth and recipient site as well as factors that can be adjusted by the operator during treatment. Non-adjustable anatomical and morphological characteristics of the donor tooth and recipient site include the presence of periodontium in the donor tooth, morphology, root development stage, functionality of the donor tooth, and the presence of periodontium in the extraction socket. Factors that can be adjusted by the operator to improve the likelihood of success and survival of the transplanted tooth include reduction of extra-oral exposure time of the donor tooth, tight contact between the donor tooth and recipient site, 4 - 8 weeks of non-rigid splinting to allow physiological mobility of the donor tooth, and the application of bite force during the initial post-treatment period to prevent ankylosis[10,11]. The 2 most influential factors in the autotransplantation of teeth are precise shaping of the recipient site, and reduction in the operating time to minimize damage to the periodontal membrane of the donor tooth[10,11].

For autotransplantations performed in previous studies, only rapid prototyping (RP) models of the donor tooth and recipient site were manufactured. This aimed to prevent periodontal damage to the donor tooth caused by repeatedly checking of the fit of the donor tooth in the extraction socket[10,11]. However, as only RP models were used, the extraction sockets underwent excessive removal, and the treatment time was relatively long. Furthermore, treatment precision was reduced due to differences in size between the manufactured models and the corresponding donor tooth and recipient site[10,11].

To overcome these limitations, in the present study, more advanced than the computer-aided rapid prototyping (CARP) models of donor tooth and recipient site, additional 3-dimensional (3D)-image analysis and 3D-printing technology were used. These advancements are gaining popularity in contemporary dental fields of prosthodontics, orthodontics, and oral

surgery[12,13]. After casting models of the donor tooth and recipient site, surgical guide templates, which can guide the direction and depth of bone removal to prevent excessive removal of the extraction socket, were created using 3D-printing technology. This was followed by metal casting of the templates for bone fracture resistance during bone removal. These manufactured models and surgical guide templates reduce the time required for recipient site formation compared with previous methods. Furthermore, damage to the periodontium of the donor tooth is decreased, as the donor tooth does not need to be checked multiple times to ensure that it fits into the recipient site.

The present case of autotransplantation of a tooth with complete roots at the site of a congenitally missing permanent tooth using surgical guide templates generated with 3D-printing technology and donor tooth and recipient site models yielded favorable results.

II. Case Report

An 18-year-old girl visited the Department of Pediatric Dentistry of OO Dental Hospital for transplanting the right maxillary 3rd molar to the site of a right mandibular 2nd premolar, which is congenitally missing, before orthodontic treatment. A congenitally missing right mandibular 2nd premolar, dental caries of the right mandibular 2nd primary molar, and a sound right maxillary 3rd molar were observed on a panoramic radiograph (Fig. 1).

We performed post-transplantation analysis using Invivo™ 5 (Anatomage Inc., USA) and Geomagic® Design™ X software (3D Systems, USA), based on cone beam computed tomography (CBCT) images.




Fig. 1. Panoramic radiographic view at initial visit shows a congenitally missing right mandibular 2nd premolar, dental caries of the right mandibular 2nd primary molar, and a sound right maxillary 3rd molar.




Fig. 2. Analysis of proximal space with transplantation of the right maxillary 3rd molar into the site of the right mandibular 2nd primary molar as it is shaped (lingual view).




Fig. 3. Analysis of proximal space with transplantation of the right maxillary 3rd molar into the site of the right mandibular 2nd primary molar with 90° rotation (lingual view).

First, we analyzed the proximal space of the transplanted site by transplanting the right maxillary 3rd molar into the site of the right mandibular 2nd primary molar. The results showed that if the right maxillary 3rd molar were transplanted as it was shaped, a small amount of proximal space would remain, and approximately half of the lingual root surface of the right maxillary 3rd molar would be exposed (Fig. 2). By contrast, transplantation of the right maxillary 3rd molar into the site of the right mandibular 2nd primary molar with 90° rotation would leave no proximal space and the lingual root surface exposure would be reduced to approximately half that associated with transplantation without rotation (Fig. 3).

Additionally, transplantation of the right maxillary 3rd molar with 90° rotation, there would provide sufficient distance from the mental foramen (Fig. 4). With regard to the comparison of the extraction socket width of the right mandibular 2nd primary molar with the root width of the right maxillary 3rd molar, the extraction socket was slightly larger, but similar enough to render the transplantation viable (Fig. 5).




Fig. 4. Analysis of the distance from the mental foramen to the root apex of the transplanted teeth.




Fig. 5. Comparison of the extraction socket of the right mandibular 2nd primary molar and the root width of the right maxillary 3rd molar.

Furthermore, transplantation of the right maxillary 3rd molar vertically into the extraction socket would necessitate removal of buccal bone from the extraction socket. However, although transplantation of the donor tooth with buccal tipping would increase the lingual root surface exposure, it would also decrease alveolar bone removal. For this reason, we planned transplantation with buccal tipping (Fig. 6).

Based on the results of these analyses, to properly transplant the right maxillary 3rd molar into the extraction socket, we manufactured 3 surgical guide templates using 3D-printing technology.

The 1st guide template was designed to prevent damage to the periodontium by replicating the donor tooth and fitting it into the recipient site repeatedly. The 2nd guide template was needed to ensure the correct depth of bone removal at the extraction socket. The 3rd guide template was designed to prevent excessive bone removal in the bucco-lingual direction at the recipient site (Fig. 7).

With these surgical guide templates, we performed auto-




Fig. 6. Comparison of the major axis of the donor tooth and bone removal requirement at the recipient site according to direction of transplantation. (A) Transplanted vertically. (B) Transplanted with buccal tipping.




Fig. 7. Surgical guide templates created using 3D-printing technology. (A) Replicas of donor tooth. (B) Guiding the correct depth of the recipient site. (C) Guiding the correct bucco-lingual direction of the recipient site.

transplantation of the right maxillary 3rd molar into the site of the right mandibular 2nd primary molar. First, the right mandibular 2nd primary molar at the recipient site was extracted by sectioning it in half (Figs. 8A and 8B). The required depth of the extraction socket was achieved using the surgical guide templates and implant burs, followed by the creation of the recipient site using a conventional round surgical bur (Fig. 8C).

The right maxillary 3rd molar was carefully extracted with minimal damage to the periodontium and immediately stored in saline (Fig. 8D). The root apex of the donor tooth was removed and retrofilled with mineral trioxide aggregate (Fig. 8E). The donor tooth was then transplanted into the prepared recipient site without any contact with the antagonistic tooth, a suture splint was then applied (Fig. 8F). After 1 week, we decided that it would be difficult to stabilize the donor tooth with suturing alone, so the tooth was splinted using a multi-twist wire (Fig. 8G). The splinted wire was removed after 3 weeks. 4 weeks




Fig. 8. Treatment procedures. (A), (B) The right mandibular 2nd primary molar at the recipient site is extracted after sectioning it in half. (C) The recipient site is formed using the surgical guide templates and implant burs. (D) The donor tooth is extracted carefully. (E) The root apex of the donor tooth is cut and retrofilled with mineral trioxide aggregate. (F) The donor tooth is transplanted to the recipient site and splinted via sutures. (G) After 1 week, a multi-twist wire splint is placed for 3 weeks. (H) 4 weeks after the transplantation, endodontic treatment with mineral trioxide aggregate is performed.




Fig. 9. Healing state from transplantation of the donor tooth to post-transplantation. (A) At 4 weeks, (B) at 2 months, (C) at 3 months, and (D) at 6 months.

after the transplantation, endodontic treatment was performed using mineral trioxide aggregate (Fig. 8H). And the patient is currently undergoing orthodontic treatment.

Gradual bone formation was observed around the transplanted tooth at 4 weeks, 2 months, and 3 months post-transplantation. And at 6 months post-transplantation, sufficient bone had formed around the periodontium of the transplanted tooth, and the alveolar bone had developed a favorable trabecular pattern. There were no complications, such as ankylosis, inflammatory root resorption, or periapical lesions (Fig. 9).

III. Discussion

Autotransplantation is commonly used for tooth replacement procedures, especially in adolescents, since implants and other prosthetic replacements are contraindicated for various reasons[14-16].

During surgery, the operator can adjust some of the parameters that affect the prognosis of autotransplantation such as extra-oral time and micro-movement of the donor tooth.

A previous study found that most of the periodontium survived when left dry for 18 minutes extraorally. However, when left dry for 30 minutes, more than half died, and when left dry for 120 minutes, the majority died[17]. Therefore, in cases of

autotransplantation, it is recommended that extra-oral time of the donor tooth should not exceed 30 minutes.

Transplantation of a donor tooth into a recipient site that is too large can slow bone healing. Also, if the distance between the donor tooth and recipient site is too narrow, ankylosis can occur on the damaged periodontium[18]. Therefore, the distance between the donor tooth and the recipient site must be appropriate to permit micro-movement of the tooth.

Previous studies of autotransplantation used only CARP models of the donor tooth and recipient site. However, in the present case, 3 additional surgical guide templates were created using 3D-printing technology. The most commonly reported complications associated with autotransplanted teeth are ankylosis and inflammatory root resorption[7,18]. Survival of the periodontium of the donor tooth is essential to prevent these complications[7,18]. As in the present case, the use of multiple surgical guide templates can reduce periodontium damage and increase the success and survival rates of autotransplantation by facilitating a faster and safer procedure.

The surgical guide templates used in the present case were able to guide the correct bucco-lingual direction and depth of bone removal at the recipient site. However, templates reproducing bone removal at the recipient site similar to the root shape of the donor tooth could not be manufactured. More

satisfactory results may have been achieved if these additional templates had been available.

Additionally, in the present case, the donor tooth was transplanted with buccal tipping to reduce bone removal, despite the increased exposure of the lingual root surface. The likelihood of favorable healing in such a case is doubtful. Transplantation of a tooth with root exposure at a site with insufficient bone yields a certain amount of bone formation, but osteogenesis up to the gingival line can be difficult to achieve[19]. However, transplantation of the donor tooth sufficiently deep to prevent exposure of the lingual root surface increases bone removal at the recipient site and ultimately leads to resorption of the alveolar bone ridge. Therefore, while the donor tooth in the present case was transplanted at an appropriate depth with some degree of lingual root exposure, a more satisfactory result may have been achieved if bone grafting had been performed on the lingual side.

The donor tooth used in the present case was a hypo-functional maxillary 3rd molar with an atrophied periodontal ligament that lacked matrix proteins and had poor blood circulation. Therefore, the most likely potential complication was ankylosis[7]. To increase the mobility of a hypo-functional tooth and volume of the periodontal ligament cells, pre-operative orthodontic extrusion can be performed for 2 - 3 weeks prior to surgery[7].

The routine autotransplantation procedure of a tooth with complete roots comprises 3 stages; autotransplantation, endodontic treatment 2 - 4 weeks after surgery, and removal of the splint 4 - 8 weeks postoperatively[7].

Andreasen *et al.*[19] reported that when a donor tooth with incomplete root formation was transplanted, the vitality of the pulp was 76 - 94%, whereas when a donor tooth with complete root formation was transplanted, it was much lower at 0 - 22%. This suggests that the conventional endodontic treatment should be performed for teeth with complete roots[20]. Additionally, when transplanting teeth with complete roots, endodontic treatment must be performed within 3 - 4 weeks after transplantation[20].

Finally, to reduce the occurrence of post-treatment ankylosis, semi-rigid splinting may be considered. This splinting method allows physiologic movement of the donor tooth[7,21]. With regard to splinting methods, Kahnberg[22] reported that some freedom of mobility of the donor tooth during the healing period can prevent ankylosis. Kristerson and Andreasen[23] found that rigid splinting increases root resorption, therefore,

allowing slight mobility of the donor tooth promotes the recovery of initial osseous ankylosis by functionally stimulating the periodontal ligament. Non-rigid splinting, such as suturing, for a short period of 1 - 2 weeks is recommended.

Additionally, early application of excessive force to the donor tooth can cause severe root resorption of the tooth and alveolar bone resorption[7]. Therefore, in the present case, the donor tooth was carefully inserted into the socket without occlusal contact to prevent excessive occlusal stimulation. A non-rigid splint of sutures was applied during the 1st week after transplantation, followed by a semi-rigid splinting with wire for 3 weeks. Furthermore, we attempted a new method of semi-rigid splinting by splinting both the buccal and lingual surfaces to the adjacent tooth with the same force, distributing the force to the tooth and wire. This resulted in a greater force on the tooth, with a lower elastic modulus than conventional splints, and allowed more movement of the donor tooth. However, further studies are needed to evaluate the physiologic mobility of teeth with this new splinting method.

The use of 3D-printing technology is becoming more prevalent in the fields of prosthodontics, oral surgery, and orthodontics, helping to improve the accuracy and safety of many treatments[12,13]. In pediatric dentistry, 3D-printing is used not only for autotransplantation, but also for proper crown setting, decreasing the time spent fabricating celluloid crowns or zirconia crowns that fit each tooth[24].

However, 3D-printing technology is not yet commonly used due to its high cost and compatibility issues, therefore, further studies and developments are required.

IV. Summary

The use of 3D-printing technology, and the fabrication of surgical guide templates in addition to models of the donor tooth and recipient site may reduce periodontal damage to the donor tooth, decrease treatment time, and increase the safety and success rate of autotransplantation.

Also, to reduce post-treatment complications such as ankylosis, in the present case, we attempted a new approach of semi-rigid splinting.

To date, the transplanted tooth has remained stable and has not developed any complications. However, because the donor teeth have a high incidence of complications such as ankylosis, regular follow-ups are required.

References

1. Aoyama S, Yoshizawa M, Saito C, *et al.* : Prognostic factors for autotransplantation of teeth with complete root formation. *Oral Surg Oral Med Oral Pathol Oral Radiol*, 114:216-228, 2012.
2. Watanabe Y, Mohri T, Saito I, *et al.* : Long-term observation of autotransplanted teeth with complete formation in orthodontic patients. *Am J Orthod Dentofacial Orthop*, 138:720-726, 2010.
3. Anssari Moin D, Derksen W, Wismeijer D, *et al.* : A Novel Approach for Computer-Assisted Template-Guided Autotransplantation of Teeth With Custom 3D Designed/Printed Surgical Tooling. An Ex Vivo Proof of Concept. *J Oral Maxillofac Surg*, 74:895-902, 2016.
4. Clokie CM, Yau DM, Chano L : Autogenous tooth transplantation: an alternative to dental implant placement?. *J Can Dent Assoc*, 67:92-96, 2001.
5. Moorrees CF, Fanning EA, Hunt EE Jr : Age variation of formation stage for ten permanent teeth. *J Dent Res*, 42:1490-1502, 1963.
6. Czochrowska EM, Stenvik A, Album B, Zachrisson BU : Autotransplantation of premolars to replace maxillary incisors: A comparison with natural incisors. *Am J Orthod Dentofacial Orthop*, 118:592-600, 2000.
7. Kokai S, Kanno Z, Soma K, *et al.* : Retrospective study of 100 autotransplanted teeth with complete root formation and subsequent orthodontic treatment. *Am J Orthod Dentofacial Orthop*, 148:982-989, 2015.
8. Yoshino K, Kariya N, Matsukubo T, *et al.* : A retrospective survey of autotransplantation of teeth in dental clinics. *J Oral Rehabil*, 39:37-43, 2012.
9. Czochrowska EM, Stenvik A, Bjercke B, Zachrisson BU : Outcome of tooth transplantation; survival and success rates 17-41 years posttreatment. *Am J Orthod Dentofacial Orthop*, 121:110-119, 2002.
10. Lee SJ, Jung IY, Kum KY, *et al.* : Clinical application of computer-aided rapid prototyping for tooth transplantation. *Dent Traumatol*, 17:114-119, 2001.
11. Lee SJ, Kim ES : Minimizing the extra-oral time in autogenous tooth transplantation; use of computer-aided rapid prototyping (CARP) as a duplicate model tooth. *Restor Dent Endod*, 37:136-141, 2012.
12. Strub JR, Rekow ED, Witkowski S : Computer-aided design and fabrication of dental restorations: Current systems and future possibilities. *J Am Dent Assoc*, 137:1289-1296, 2006.
13. Kwon SY, Kim Y, Kim SH, *et al.* : Computer-Aided Designing and Manufacturing of Lingual Fixed Orthodontic Appliance Using 2D/3D Registration Software and Rapid Prototyping. *Int J Dent*, 1-8, 2014.
14. Kim ES, Jung JY, Lee SJ, *et al.* : Evaluation of the prognosis and causes of failure in 182 cases of autogenous tooth transplantation. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 100:112-119, 2005.
15. Park JM, Tatad JC, Kim SJ, *et al.* : Optimizing Third Molar Auto-transplantation; Applications of Reverse-Engineered Surgical Templates and Rapid Prototyping of Three-Dimensional Teeth. *J Oral Maxillofac Surg*, 72:1653-1659, 2014.
16. Tsukiboshi M : Autotransplantation of teeth: requirements for predictable success. *Dent Traumatol*, 18:157-180, 2002.
17. Andreasen JO : Effect of extra-alveolar period and storage media upon periodontal and pulpal healing after replantation of mature permanent incisors in monkeys. *Int J Oral Surg*, 10:43-53, 1981.
18. Jang YJ, Choi YJ, Kim ES, *et al.* : Prognostic Factors for Clinical Outcomes in Autotransplantation of Teeth with Complete Root Formation: Survival Analysis for up to 12 Years. *J Endod*, 42:198-205, 2016.
19. Andreasen JO, Hjorting-Hansen E, Jolst O : A clinical and radiographic study of 76 autotransplanted third molars. *Scand J Dent Res*, 78:512-523, 1970.
20. Mejàre B, Wannfors K, Jansson L : A prospective study on transplantation of third molars with complete root formation. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 97:231-238, 2004.
21. Bauss O, Schwestka-Polly R, Schilke R, Kiliaridis S : Effect of different splinting methods and fixation periods on root development of autotransplanted immature third molars. *J Oral Maxillofac Surg*, 63:304-310, 2005.
22. Kahnberg KE : Autotransplantation of teeth (I). Indications for transplantation with a follow-up of 51 cases. *Int J Oral Maxillofac Surg*, 16:577-585, 1987.
23. Kristerson L, Andreasen JO : The effect of splinting upon periodontal and pulpal healing after autotransplantation of mature and immature permanent incisors in monkey. *Int J Oral Surg*, 12:239-249, 1983.
24. Lee SH : Prospect for 3D Printing Technology in Medical, Dental, and Pediatric Dental Field. *J Korean Acad Pediatr Dent*, 43:93-108, 2016.

국문초록

3D 프린팅 기술을 이용한 치근단 완성 치아의 자가이식에 대한 최신 접근

박정하¹ · 이상호¹ · 이난영¹ · 지명관¹ · 정혜란²

¹조선대학교 치의학전문대학원 소아치과학교실

²동아보건대학교 치위생과

교정치료의 목적으로, 또는 보존 불가능한 치아를 발거 후 치근 형성이 완료된 치아의 자가치아이식을 요구하는 경우가 늘어나고 있지만 미완성 치근 형성 치아를 이식하는 경우에 비하여 치근 형성이 완료된 치아를 이식하는 경우 생존률 및 성공률이 낮음이 보고되고 있다.

이전의 자가치아이식에서는 공여치와 수여부만을 재현하는 모델만이 사용된 반면에 현재는 불필요한 골삭제와 술식 시간을 감소시켜주는 외과적 가이드 템플레이트들을 3D 프린팅 기술을 통하여 제작, 술식에 추가적으로 이용함으로써 이식시 적절한 방향과 깊이로 수여부를 형성할 수 있으며 술식시간도 감소시켜 자가치아이식의 성공률 및 생존률을 향상시킬 수 있다고 할 수 있다.

본 케이스는 치근 형성이 완료된 치아를 선천성 결손 부위에 자가이식한 케이스로 3D 프린팅 기술을 이용하여 공여치와 수여부 모형 모델 및 외과적 가이드 템플레이트들을 제작함으로써 술식의 성공률과 안정성을 높일 수 있었고 이식치의 양호한 치유 결과를 이 끌어냈다.